

Post– Emergent Weed Control in Faba Beans—2018

Herbicide	Herbicide Group	Application Stage*	Tank-Mix Options (supported or on label)	Primary Target	Re-cropping (year after)
Basagran®/Basagran® Forte/ Benta Super/Broadloom® (bentazon)	6	From 2 leaf stage or 4 inches in height and larger	None on label	Broadleaf weeds	None
Select®/Centurion®/Shadow® RTM (clethodim)	1	Any (60 day pre-harvest interval)	None on label	Grassy weeds	None
Odyssey® NXT/Duet™ (imaxamox + imazethapyr)	2	1 to 6 leaf (1 to 6 above-ground)	None on label	Broadleaf weeds Grassy weeds	Crop restrictions on label
Odyssey® Ultra NXT (imaxamox + mazethapyr + sethoxymdim)	1 + 2	1 to 6 leaf (1 to 6 above-ground node)	Poast® Ultra	Broadleaf weeds Grassy weeds	Crop restrictions on label
Poast® Ultra (sethoxymdim)	1	Any (80 day pre-harvest interval)	Odyssey® NXT	Grassy weeds	None
Assure® II/Yuma™ GL/ Contender® (quizalofop)	1	Any (30 day pre-harvest interval)	Odyssey® NXT	Grassy weeds	None
Viper® ADV (Imazamox + bentazon)	2 + 6	1 to 6 leaf (1 to 6 above-ground)	None on label	Broadleaf weeds Grassy weeds	Crop restrictions on label

*Application stages on labels may refer to leaf, node, or above-ground node stages.

There may be restrictions on label for rates, varieties, or soil types. Consult label or manufacturer for more detail. Always read and follow label directions. This information is only a guide, if label information differs, follow label instructions.