

## Field Pea Fungicide Options 2018

Fungicide	Fungicide Group	Maximum Number of Applications Registered Per Year**	Aerial Application	Pre-Harvest Interval	Diseases (Pathogens)									
					Anthraxnose ( <i>Colletotrichum</i> spp.)	Ascochyta Complex ( <i>Ascochyta</i> spp.)	Ascochyta Complex ( <i>Mycosphaerella pinodes</i> )	Brown Spot ( <i>Pseudomonas syringae</i> )	Downy Mildew ( <i>Peronospora viciae</i> )	Grey Mould ( <i>Botrytis cinerea</i> )	Powdery Mildew ( <i>Erysiphe pisi</i> )	Root Rot ( <i>Aphanomyces euteiches</i> , <i>Pythium ultimum</i> )	Sclerotinia ( <i>Sclerotinia sclerotiorum</i> )	
<b>Acapela®</b> (picoxystrobin)	11	2	Yes	14 days			X*							X*
<b>Quadris®/Azoshy 250SC</b> (azoxystrobin)	11	2	Yes	15 days	X	X	X				X			X*
<b>Bravo®/Echo</b> (chlorothalonil)	M5	3	Yes	32 days			X							
<b>Cueva®</b> (copper)	M1	Contact manufacturer	No	1 day		X		X			X			
<b>Cosavet DF Edge</b> (sulphur)	M2	8	No	1 day							X			
<b>Cotegra™</b> (boscalid + prothioconazole)	3 + 7	2	Yes	21 days										X*
<b>Delaro™</b> (prothioconazole + trifloxystrobin)	3 + 11	2	Yes	30 days		X	X			X				X
<b>Dyax™</b> (fluxapyroxad + pyraclostrobin)	7 + 11	2	Yes	30 days			X				X			X*
<b>Elatus™</b> (azoxystrobin + benzovindiflupyr)	7 + 11	1	Yes	15 days	X	X	X				X			X*
<b>Headline® EC</b> (pyraclostrobin)	11	2	Yes	30 days		X	X		X		X			
<b>Kenja® 400SC</b> (isofetamid)	7	2	No	30 days										X*
<b>Lance® AG</b> (boscalid + pyraclostrobin)	7 + 11	2	Yes	30 days		X	X		X*	X	X			

Fungicide	Fungicide Group	Maximum Number of Applications Registered Per Year**	Aerial Application	Pre-Harvest Interval	Diseases (Pathogens)								
					Anthraco- nose ( <i>Colletotrichum</i> spp.)	Ascochyta Complex ( <i>Ascochyta</i> spp.)	Ascochyta Complex ( <i>Mycosphaerella pinodes</i> )	Brown Spot ( <i>Pseudomonas syringae</i> )	Downy Mildew ( <i>Peronospora viciae</i> )	Grey Mould ( <i>Botrytis cinerea</i> )	Powdery Mildew ( <i>Erysiphe pisi</i> )	Root Rot ( <i>Aphanomyces euteiches</i> , <i>Pythium ultimum</i> )	Sclerotinia ( <i>Sclerotinia sclerotiorum</i> )
<b>Lance® WDG</b> (boscalid)	7	2	No	21 days		X	X			X			
<b>Phostrol™</b> (phosphites)	33	2	No	21 days								X*	
<b>Priaxor®</b> (fluxapyroxad + pyralostrobin)	7 + 11	2	Yes	21 days			X		X*		X		X*
<b>Tilt®/Propel®/ Propi Super 25EC</b> (propiconazole)	3	2	Yes	30 days							X		
<b>Quash®</b> (metconazole)	3	2	Yes	21 days							X		X*
<b>Quilt®/Fungtion SC</b> (azoxystrobin +	3 + 11	2	Yes	30 days			X				X		
<b>Serenade®</b> ( <i>Bacillus subtilis</i> QST 713 strain)	44	None	No	None						X			X

\* Product registered for suppression only.

\*\* Follow label for proper rotation of fungicides with different modes of action. Some products are not labelled for two sequential applications. Rates and staging may vary by fungicide and disease pressure. Read and follow label directions.