

## Post-Emergent Weed Control in Chickpeas

Herbicide**	Herbicide Group	Application Stage*	Tank-Mix Options (on label)	Primary Target	Re-cropping (year after)
<b>Select®/Centurion®/ Statue™/ Arrow®/Arrow® All-in/Shadow™ RTM/Patron 240/Antler™/ MPower® Independence™</b> (clethodim)	1	Up to the ninth above-ground node stage, and no larger than seven inches in height	None on label	Grassy weeds	None
<b>Sencor®/TriCor®/Squadron®/ Squadron/Buzzin® 70WDG/ Metrix SC</b> (metribuzin)	5	Up to 2.5 inches in height when vines have one to three above-ground nodes	None on label	Broadleaf weeds	Crop restrictions on label
<b>Solo® ADV**</b> (imaxamox)	2	One to six above-ground node	None on label	Broadleaf weeds Grassy weeds	Crop restrictions on label
<b>Poast® Ultra</b> (sethoxydim)	1	Any (65 day pre-harvest interval)	None on label	Grassy weeds	None
<b>Assure® II/Yuma™ GL/Contender®</b> (quizalofop)	1	Any (85 day pre-harvest interval)	None on label	Grassy weeds	None

\*Application stages on labels may refer to leaf, node, or above-ground node stages.

\*\*Herbicide is for use on imidazolinone tolerant chickpea varieties only.

There may be restrictions on label for rates, varieties, or soil types. Consult label or manufacturer for more detail. Always read and follow label directions. This information is only a guide, if label information differs, follow label instructions.

Version March 2019