

Post-Emergent Weed Control in Dry Beans

Herbicide	Herbicide Group	Application Stage	Tank-Mix Options (supported or on label)	Primary Target	Re-cropping (year after)
Basagran®/Basagran® Forte/ Benta Super/Broadloom® (bentazon)	6	After the first trifoliolate leaf	None on label	Broadleaf weeds	None
Select®/Centurion®/Statue®/ Arrow®/Arrow® All-in /Shadow™ RTM/Patron 240/Antler™/MPower® Independence™ (clethodim)	1	Any (60 day preharvest interval) Not all varieties have been tested for tolerance Check label for varieties	Pursuit®	Grassy weeds	None
Pursuit®/MPower®/Kamikaze®/ Phantom®/Guardman® Gladiator™/Multistar® (imazethapyr)	2	Up to, and including, the second trifoliolate	None on label	Broadleaf weeds Grassy weeds	Crop restrictions on label
Permit® (halosulfuron)	2	Second to fourth trifoliolate (prior to flower) Not all varieties have been tested for tolerance Check label for varieties	Yuma™ GL	Broadleaf weeds	Crop restrictions on label
Poast® Ultra (sethoxydim)	1	Any (80 day pre-harvest interval)	None on label	Grassy weeds	None
Assure® II/Yuma™ GL/Contender® (quizalofop)	1	Any (65 day pre-harvest interval)	Basagran®	Grassy weeds	None
Viper® ADV (Imazamox + bentazon)	2 + 6	First fully expanded trifoliolate to second fully expanded trifoliolate Not all varieties have been tested for tolerance Check label for varieties	Basagran® Forte	Broadleaf weeds Grassy weeds	Crop restrictions on label

There may be restrictions on label for rates, varieties, or soil types. Consult label or manufacturer for more detail. Always read and follow label directions. This information is only a guide, if label information differs, follow label instructions.

Version March 2019