

Faba Bean Fungicide Options 2019

Fungicide	Fungicide Group	Maximum Number of Applications Registered Per Year**	Aerial Application	Pre-Harvest Interval	Diseases (Pathogens)				
					Ascochyta Blight (Ascochyta spp.)	Anthraxnose (Colletotrichum spp.)	Grey mold (Botrytis cinerea)	Powdery Mildew (Erysiphe spp.)	Sclerotinia
Acapela® (picoxystrobin)	11	2	Yes	14 days					X*
Quadris®/Azoshy 250 SC (azoxystrobin)	11	2	Yes	15 days	X	X			X*
Cotegra™ (boscalid + prothioconazole)	3 + 7	2	Yes	21 days					X*
Dyax™ (fluxapyroxad + pyraclostrobin)	7 + 11	2	Yes	30 days	X*				X*
Elatus™ (azoxystrobin + benzovindiflupyr)	7 + 11	1	Yes	15 days (Apply no later than 20 per cent bloom)	X	X			X*
Headline® EC (pyraclostrobin)	11	2	Yes	30 days	X			X	
Kenja™ 400SC (isofetamid)	7	2	No	30 days					X*
Lance® WDG (boscalid)	7	2	Yes	21 days					X
Priaxor® (fluxapyroxad + pyraclostrobin)	7 + 11	2	Yes	30 days	X		X*	X	X*
Propulse® (fluopyram + prothioconazole)	3 + 7	2	No	14 days	X	X			X
Tilt 250E® / Propel® / Propi Super 25EC (propiconazole)	3	2	Yes	30 days				X	
Quilt®/Fungtion SC (azoxystrobin + propiconazole)	3 + 11	2	Yes	30 days				X	

* Product registered for suppression only.

** Follow label for proper rotation of fungicides with different modes of action. Some products are not labelled for two sequential applications. Rates and staging may vary by fungicide and disease pressure. Read and follow label directions.