

## Lentil Fungicide Options 2019

Fungicide	Fungicide Group	Maximum Number of Applications Registered Per Year**	Aerial Application	Pre-Harvest Interval	Diseases (Pathogens)					
					Anthracnose ( <i>Colletotrichum truncatum</i> )	Ascochyta Blight ( <i>Ascochyta lentis</i> )	Brown Spot ( <i>Pseudomonas syringae</i> )	Grey Mould ( <i>Botrytis cinerea</i> )	Powdery Mildew ( <i>Erysiphe spp</i> )	Sclerotinia ( <i>Sclerotinia sclerotiorum</i> )
<b>Acapela®</b> (picoxystrobin)	11	2	Yes	14 days	X	X				X*
<b>Quadris®/Azoshy 250 SC</b> (azoxystrobin)	11	2	Yes	15 days	X	X				X*
<b>Bravo 500 ZN®/Echo 720</b> (chlorothalonil)	M5	2	Yes	48 days	X	X				
<b>Cueva®</b> (copper)	M1	15	No	1 day			X		X	
<b>Cotegra™</b> (boscalid + prothioconazole)	3 + 7	2	Yes	21 days						X*
<b>Delaro 325 SC™</b> (prothioconazole + trifloxystrobin)	3 + 11	2	Yes	30 days	X	X		X		X
<b>Dithane™ Rainshield/ Manzate® Pro-Stick/ Penncozeb 75DF®</b> (Mancozeb)	M3	3	Yes	35 days	X	X				
<b>Dyax™</b> (fluxapyroxad + pyraclostrobin)	7 + 11	2	Yes	30 days	X	X				X*
<b>Elatus™</b> (azoxystrobin + benzovindiflupyr)	7 + 11	1	Yes	15 days (Apply no later than 20 per cent bloom)	X	X				X*
<b>Headline® EC</b> (pyraclostrobin)	11	2	Yes	30 days	X	X				
<b>Kenja™ 400SC</b> (isofetamid)	7	2	No	30 days						X*

Fungicide	Fungicide Group	Maximum Number of Applications Registered Per Year**	Aerial Application	Pre-Harvest Interval	Diseases (Pathogens)					
					Anthraxnose ( <i>Colletotrichum truncatum</i> )	Ascochyta Blight ( <i>Ascochyta lentis</i> )	Brown Spot ( <i>Pseudomonas syringae</i> )	Grey Mould ( <i>Botrytis cinerea</i> )	Powdery Mildew ( <i>Erysiphe spp.</i> )	Sclerotinia ( <i>Sclerotinia sclerotiorum</i> )
<b>Lance® AG</b> (boscalid + pyraclostrobin)	7 + 11	2	Yes	30 days	X	X		X		X
<b>Lance® WDG</b> (boscalid)	7	2	Yes	21 days		X		X		X
<b>Priaxor®</b> (fluxapyroxad + pyraclostrobin)	7 + 11	2	Yes	30 days	X	X		X*		X*
<b>Proline® 480 SC</b> (prothioconazole)	3	2	Yes	7 days		X				X
<b>Tilt 250E®/Propel®/ Propi Super 25 EC</b> (propiconazole)	3	2	Yes	30 days					X	
<b>Quash®</b> (metconazole)	3	2	Yes	21 days		X*				X*
<b>Quilt®/Fungtion SC</b> (azoxystrobin + propiconazole)	3 + 11	2	Yes	30 days	X				X	
<b>Serenade OPTI®</b> ( <i>Bacillus subtilis</i> QST 713 strain)	44	None	No	None				X		X
<b>Topnotch®</b> (azoxystrobin + propiconazole)	3 + 11	2	Yes	30	X	X			X	X*

\* Product registered for suppression only.  
\*\* Follow label for proper rotation of fungicides with different modes of action. Some products are not labelled for two sequential applications. Rates and staging may vary by fungicide and disease pressure. Read and follow label directions.