

## Soybean Fungicide Options 2019

Fungicide	Fungicide Group	Maximum Number of Applications Registered Per Year**	Aerial Application	Pre-Harvest Interval	Diseases (Pathogens)									
					Anthracnose ( <i>Colletotrichum truncatum</i> )	Bacterial Blight ( <i>Pseudomonas syringae</i> )	Brown Spot ( <i>Septoria glycines</i> )	Cercospora Leaf Spot ( <i>Cercospora kikuchii</i> )	Frogeye Leafspot ( <i>Cercospora soja</i> )	Phomopsis Stem Blight ( <i>Phomopsis longicolla</i> )	Pod and Stem Blight ( <i>Diaporthe phaseolorum</i> )	Powdery Mildew ( <i>Erysiphe</i> spp.)	White Mold ( <i>Sclerotinia sclerotiorum</i> )	
<b>Acapela®</b> (picoxystrobin)	11	2	Yes	14 days			X		X					X*
<b>Allegro® 500F</b> (fluazinam)	29	2	Yes	Do not apply after R3 (early pod)										X
<b>Quadris®/Azoshy 250SC</b> (azoxystrobin)	11	2	Yes	15 days				X				X		
<b>Contans® WG</b> ( <i>Coniothyrium minitans</i> )	Not classified Bio-fungicide	Pre-Plant or post-harvest	No	None										X
<b>Cueva®</b> (copper)	M1	15	No	1 day		X						X		
<b>Cotegra™</b> (boscalid +)	3 + 7	2	Yes	21 days			X*		X					X*
<b>Delaro™</b> (prothioconazole + trifloxystrobin)	3 + 11	2	Yes	20 days			X		X	X				X
<b>Double Nickel™ LC/Double Nickel™ 55</b> ( <i>Bacillus amyloliquefaciens</i> strain D747)	44	Every three to 10 days	No	None										X
<b>Dyax™</b> (fluxapyroxad + pyraclostrobin)	7 + 11	2	Yes	21 days			X*		X*					X*
<b>Elatus</b> (azoxystrobin + benzovindiflupyr)	7 + 11	2	Yes	14 days			X	X	X*		X*	X		

Fungicide	Fungicide Group	Maximum Number of Applications Registered Per Year**	Aerial Application	Pre-Harvest Interval	Diseases (Pathogens)								
					Anthraxnose ( <i>Colletotrichum truncatum</i> )	Ascochyta Blight ( <i>Ascochyta</i> spp.)	Brown Spot ( <i>Septoria glycines</i> )	Cercospora Leaf Spot ( <i>Cercospora kikuchii</i> )	Frogeye Leafspot ( <i>Cercospora sojina</i> )	Phomopsis Stem Blight ( <i>Phomopsis longicolla</i> )	Pod and Stem Blight ( <i>Diaporthe phaseolorum</i> )	Powdery Mildew ( <i>Erysiphe</i> spp.)	White Mold ( <i>Sclerotinia sclerotiorum</i> )
<b>Fullback™ 125SC</b> (flutriafol)	3	3	No	21 days			X	X					
<b>Headline® EC</b> (pyraclostrobin)	11	2	Yes	21 days					X				
<b>Priaxor®</b> (fluxapyroxad + pyraclostrobin)	7 + 11	2	Yes	21 days			X						X*
<b>Proline®</b> (prothioconazole)	3	1	Yes	20 days					X				
<b>Tilt®/Propel®/Propi Super 25EC/ Bumper 432 EC®/ Pivot 418 C®/ FITNESS®/Nufarm Propiconazole Fungicide / Co-Op Pivot</b> (propiconazole)	3	2	Yes	30 –50 days, depending on product used				X	X (soybean grown for seed only)			X	
<b>Quilt®/Fungtion SC</b> (azoxystrobin + propiconazole)	3 + 11	2	Yes	30 days	X				X			X	X*
<b>Serenade®</b> ( <i>Bacillus subtilis</i> )	44	None	No	None			X		X				X
<b>Folicur® 250EW</b> (tebuconazole)	3	1	Yes	36 days					X			X	
<b>Topnotch®</b> (azoxystrobin + propiconazole)	3 + 11	2	Yes	30	X							X	X*
<b>Trivapro™</b> (azoxystrobin + propiconazole + benzovindiflupyr)	3 + 7 + 11	2	No	30 days	X		X*		X		X	X	X*

\* Product registered for suppression only.

\*\* Follow label for proper rotation of fungicides with different modes of action. Some products are not labelled for two sequential applications. Rates and staging may vary by fungicide and disease pressure. Read and follow label directions.