Pulse Knowledge



Faba Bean Seed Treatment Options

Disease Reference — Action Thresholds and Registered Seed Treatment Options

| | | | Seed Treatment Options | | | | | | | | |
|--|---|--|------------------------|----------------|-----------------------------|---------------|------------------------|------------------|---------------------------|--------------------------|---------------------------------|
| Pathogen (Disease) | Possible Risk on Seed | Recommendation if Over Risk Level | Agrox® FL | Apron® Advance | ApronMaxx® RTA with INTEGO™ | Insure Pulse® | INTEGO™ Solo Fungicide | Trilex® EverGol® | Trilex® EverGol® SHIELD** | Vibrance® Maxx RTA / RFC | Vibrance® Maxx RFC with INTEGO™ |
| Aphanomyces euteiches | Not applicable Soil-borne disease only | Use seed treatment if: History of disease, provides early season protection only | | | X* | | Х* | | | | Х* |
| Ascochyta fabae | Greater than 10% | Use seed treatment | | | | Х | | X* | X* | | |
| Botrytis spp. (seed-borne only) | 10% on seed (total of Fusarium, Botrytis, and Sclerotinia) | Use seed treatment | | | | Х* | | Х | Х | | |
| Colletotrictrum truncatum (Anthracnose) | Rarely found on seed Primarily soil-borne disease | Use seed treatment if: History of disease, expecting humid/moist conditions | | Х | | X* | | | | | |
| Fusarium | 10% on seed (total of Fusarium, Botrytis, and Sclerotinia) Soil-borne as well | Use seed treatment | | Х | Х | Х | | Х | Х | Х | Х |
| Pythium | Not applicable Soil-borne disease only | Use seed treatment if: History of disease, seeding under cool-moist conditions | | Х | Х | Х | х | Х | Х | Х | Х |
| Rhizoctonia | Unknown Primarily soil-borne disease | Use seed treatment if: History of disease, seeding under cool-moist conditions | | Х | Х | Х | | Х | Х | Х | Х |
| Sclerotinia sclerotiorum (seed-borne only) *** | 10% on seed (total of Fusarium, Botrytis, and Sclerotinia) | Use seed treatment | | | | | | | | | |
| General seed rot, root rot, damping-off, and seedling | Not applicable | Use seed treatment if: History of disease, seeding under cool-moist conditions | Х | | | | | | | | |

^{*} Product registered for suppression only

(see next page for seed treatments to manage insect pests)

^{**} Product contains an insecticide component

^{***} no products registered for seed-borne $\it Sclerotinia\ sclerotiorum\ control\ in\ faba\ beans$

Faba Bean Seed Treatment Options (continued)

Insect Reference—Recommendations and Registered Seed Treatment Options

| | | Seed Treatment Options | | | | | | |
|----------------------------|--|------------------------|--------------------------------|--------------|--------------------|---------------------------|--|--|
| | | Cruiser® 5FS | CruiserMaxx® Vibrance Pulses** | Lumivia™ CPL | Stress Shield® 600 | Trilex® EverGol® SHIELD** | | |
| Insect | Recommendations Use seed treatment if: | | | | | | | |
| Armyworm and Cutworm | History of insect pressure on the field or in the nearby area | | | Х | | | | |
| Pea Leaf Weevil | Use seed treatment if: History of insect pressure on the field or in the nearby area | Х | Х | Х | Х | Х | | |
| Wireworm | Use seed treatment if: History of insect pressure on the field or in the nearby area | Х | Х | | Х | Х | | |

^{**} Product contains fungicide components

Action thresholds have not been identified for faba bean disease and insect risks, so the recommendations provided are based on thresholds from similar crops, or rules of thumb.

Understanding seed quality and using seed treatments on pulse seed when risk of disease or insect damage is present, is critical to maximizing yield.

Fungicide seed treatments protect the seed and seedlings from soil-borne diseases, as well as those diseases found on the surface of the seed or carried inside of the seed. Seed treatments will not act as a cure for seed that has high disease levels or is poor quality.

In order for a seed treatment to provide effective protection, it is important to ensure that adequate coverage of the seed is achieved and appropriate rates are used for the target pest(s).

Seed treatments will breakdown as the plant begins to develop and as a result, their limit of protection to the seed will only last for the early part of the plants growth.

When using a seed treatment in combination with an inoculant, it is important to check with the inoculant manufacturer to ensure that there is compatibility with the seed treatment, prior to use.

Seed treated with a seed treatment may pose a risk to other organisms or plants in the environment. Dispose of all leftover treated seed properly, and do not leave treated seed on the soil surface.

Version April 2020