

Chickpea Fungicide Options

| Fungicide | Fungicide Group | Maximum Number of Applications Registered Per Year** | Aerial Application | Pre-Harvest Interval | Diseases (Pathogens) | | | | | |
|---|--------------------------------|--|--------------------|----------------------------------|--|---|---|---|---|--|
| | | | | | Anthraxnose (<i>Colletotrichum truncatum</i>) | Ascochyta Blight (<i>Ascochyta lentis</i>) | Brown Spot (<i>Pseudomonas syringae</i>) | Grey Mould (<i>Botrytis cinerea</i>) | Powdery Mildew (<i>Erysiphe spp</i>) | Sclerotinia (<i>Sclerotinia sclerotiorum</i>) |
| Acapela® (picoxystrobin) | 11 | 2 | Yes | 14 days | | | | | | X* |
| Azoshy 250 SC/Quadris® (azoxystrobin) | 11 | 2 | Yes | 15 days | X | X | | | | X* |
| Bravo 500 ZN®/Echo 720 (chlorothalonil) | M5 | 2 | No | Bravo®: 14 days Echo: 48 days | | X | | | | |
| Cotegra™ (boscalid + prothioconazole) | 3 + 7 | 2 | Yes | 21 days | | | | | | X* |
| Cueva®/Copper 53W (copper) | M1 | 15 | No | 1 day | | | X | | X | |
| Delaro 325 SC™ (prothioconazole + trifloxystrobin) | 3 + 11 | 2 | Yes | 30 days | | X | | X | | X |
| Dyax™ (fluxapyroxad + pyraclostrobin) | 7 + 11 | 2 | Yes | 30 days | | X | | | | X* |
| Elatus™ (azoxystrobin + benzovindiflupyr) | 7 + 11 | 1 | Yes | 15 days | X | X | | | | X* |
| Fungtion SC/Quilt® (azoxystrobin + propiconazole) | 3 + 11 | 2 | Yes | 30 days | | | | | X | |
| Headline® EC/MPower Spade/ Preach (pyraclostrobin) | 11 (Must be mixed with Lance®) | 2 | Yes | 30 days | | X | | | | |
| Kenja™ 400SC (isofetamid) | 7 | 2 | No | 30 days | | | | | | X* |
| Lance® AG (boscalid + pyraclostrobin) | 7 + 11 | 2 | Yes | 30 days | | X | | X | | X |
| Lance® WDG (boscalid) | 7 | 2 | Yes | 21 days | | X | | X | | X |
| Miravis® Neo 300 SE (azoxystrobin + propiconazole + pydiflumetofen) | 3, 7, 11 | 1 | Yes | 30 days | X | X | | | X | X* |
| Priaxor® (fluxapyroxad + pyraclostrobin) | 7 + 11 | 2 | Yes | 30 days | | X | | | | X* |
| Proline® GOLD (fluopyram + prothioconazole) | 3 + 7 | 2 | Yes | 14 days | | X | | | | X |

(continued on next page)

Chickpea Fungicide Options (continued)

| Fungicide | Fungicide Group | Maximum Number of Applications Registered Per Year** | Aerial Application | Pre-Harvest Interval | Diseases (Pathogens) | | | | | |
|---|-----------------|--|--------------------|----------------------|--|---|---|---|---|--|
| | | | | | Anthraxnose (<i>Colletotrichum truncatum</i>) | Ascochyta Blight (<i>Ascochyta lentis</i>) | Brown Spot (<i>Pseudomonas syringae</i>) | Grey Mould (<i>Botrytis cinerea</i>) | Powdery Mildew (<i>Erysiphe spp</i>) | Sclerotinia (<i>Sclerotinia sclerotiorum</i>) |
| Proline® 480 SC (prothioconazole) | 3 | 3 | Yes | 7 days | | X | | | X | |
| Propel®/Propi Super 25EC/Tilt 250E® (propiconazole) | 3 | 2 | Yes | 30 days | | | | | | X* |
| Quash® (metconazole) | 3 | 2 | Yes | 21 days | | X* | | | X | |
| Serenade OPTI® (<i>Bacillus subtilis</i> QST 713 strain) | 44 | None | No | None | | | | X | | X |
| Zolera™ FX (fluoxastrobin + tetraconazole) | 3 + 11 | 2 | No | 14 days | | | | | | X* |

* Product registered for suppression only.

** Follow label for proper rotation of fungicides with different modes of action. Some products are not labelled for two sequential applications. Rates and staging may vary by fungicide and disease pressure. Read and follow label directions.

NOTE: Ascochyta resistance to Group 11 fungicides has been found in chickpeas. It is not recommended to apply Group 11 products alone for Ascochyta control in chickpeas.

Version March 2021