

## Faba Bean Fungicide Options

Fungicide	Fungicide Group	Maximum Number of Applications Registered Per Year**	Aerial Application	Pre-Harvest Interval	Diseases (Pathogens)				
					Ascochyta Blight (Ascochyta spp.)	Anthraxnose (Colletotrichum spp.)	Grey mold/Chocolate spot (Botrytis cinerea)	Powdery Mildew (Erysiphe spp.)	Sclerotinia
<b>Acapela®</b> (picoxystrobin)	11	2	Yes	14 days					X*
<b>Azoshy 250 SC/Quadris®</b> (azoxystrobin)	11	2	Yes	15 days	X	X			X*
<b>Cotegra™</b> (boscalid + prothioconazole)	3 + 7	2	Yes	21 days					X*
<b>Delaro 325 SC™</b> (prothioconazole + trifloxystrobin)	3 + 11	2	Yes	30 days			X		X
<b>Dyax™</b> (fluxapyroxad + pyraclostrobin)	7 + 11	2	Yes	30 days	X*				X*
<b>Elatus™</b> (azoxystrobin + benzovindiflupyr)	7 + 11	1 (apply no later than 20% bloom)	Yes	15 days	X	X			X*
<b>Fungtion SC/Quilt®</b> (azoxystrobin + propiconazole)	3 + 11	2	Yes	30 days				X	
<b>Headline® EC/MPower Spade/Preach</b> (pyraclostrobin)	11	2	Yes	30 days	X			X	
<b>Kenja™ 400SC</b> (isofetamid)	7	2	No	30 days					X*
<b>Lance® WDG</b> (boscalid)	7	2	Yes	21 days					X
<b>Miravis® Neo 300 SE</b> (azoxystrobin + propiconazole + pydiflumetofen)	3, 7, 11	1	Yes	30 days	X	X		X	X*
<b>Priaxor®</b> (fluxapyroxad + pyraclostrobin)	7 + 11	2	Yes	30 days	X		X*	X	X*
<b>Propel®/Propi Super 25EC/Tilt 250E®</b> (propiconazole)	3	2	Yes	30 days				X	
<b>Propulse®</b> (fluopyram + prothioconazole)	3 + 7	2	No	14 days	X	X			X
<b>Zolera™ FX</b> (fluoxastrobin + tetraconazole)	3 + 11	2	No	14 days		X*			X*

\*Product registered for suppression only.

\*\*Follow label for proper rotation of fungicides with different modes of action. Some products are not labelled for two sequential applications. Rates and staging may vary by fungicide and disease pressure. Read and follow label directions.

Version March 2021