

Field Pea Fungicide Options

Fungicide	Fungicide Group	Maximum Number of Applications Registered Per Year**	Aerial Application	Pre-Harvest Interval	Diseases (Pathogens)										
					Anthraxnose (Colletotrichum spp.)	Ascochyta Complex (Ascochyta spp.)	Ascochyta Complex (Mycosphaerella pinodes)	Brown Spot (Pseudomonas syringae)	Downy Mildew (Peronospora viciae)	Grey Mould (Botrytis cinerea)	Powdery Mildew (Erysiphe pisi)	Root Rot (Aphanomyces euteiches, Pythium ultimum)	Sclerotinia (Sclerotinia sclerotiorum)		
Acapela® (picoxystrobin)	11	2	Yes	14 days			X*								X*
Azoshy 250 SC/Quadris® (azoxystrobin)	11	2	Yes	15 days	X	X	X					X			X*
Bravo 500 ZN®/Echo 720 (chlorothalonil)	M5	2	Yes	32 days			X								
Cosavet DF Edge (sulphur)	M2	8	No	1 day								X			
Cotegra™ (boscalid + prothioconazole)	3 + 7	2	Yes	21 days		X*	X*								X*
Cueva® (copper)	M1	15	No	1 day		X		X				X			
Delaro 325 SC™ (prothioconazole + trifloxystrobin)	3 + 11	2	Yes	30 days		X	X			X					X
Dyax™ (fluxapyroxad + pyraclostrobin)	7 + 11	2	Yes	30 days		X	X					X			X*
Elatus™ (azoxystrobin + benzovindiflupyr)	7 + 11	2	Yes	15 days	X	X	X					X			X*
Fungtion SC/Quilt® (azoxystrobin + propiconazole)	3 + 11	2	Yes	31 days			X					X			
Headline® EC/MPower Spade/Preach (pyraclostrobin)	11	2	Yes	30 days		X	X		X			X			
Kenja® 400SC (isofetamid)	7	2	No	30 days											X*
Lance® AG (boscalid + pyraclostrobin)	7 + 11	2	Yes	30 days		X	X		X*	X	X				
Lance® WDG (boscalid)	7	2	No	21 days		X	X			X					
Miravis® Neo 300 SE (azoxystrobin + propiconazole + pydiflumetofen)	3, 7, 11	1	Yes	30 days	X	X	X					X			X*
Phostrol™ (phosphites)	33	2	No	21 days									X*		

(continued on next page)

Field Pea Fungicide Options (continued)

Fungicide	Fungicide Group	Maximum Number of Applications Registered Per Year**	Aerial Application	Pre-Harvest Interval	Diseases (Pathogens)								
					Anthraco-nose (Colletotrichum spp.)	Ascochyta Complex (Ascochyta spp.)	Ascochyta Complex (Mycosphaerella pinodes)	Brown Spot (Pseudomonas syringae)	Downy Mildew (Peronospora viciae)	Grey Mould (Botrytis cinerea)	Powdery Mildew (Erysiphe pisi)	Root Rot (Aphanomyces euteiches, Pythium ultimum)	Sclerotinia (Sclerotinia sclerotiorum)
Priaxor® (fluxapyroxad + pyralostrobin)	7 + 11	2	Yes	21 days			X		X*		X		X*
Proline® GOLD (fluopyram + prothioconazole)	3 + 7	2	Yes	14 days		X	X						X
Propel®/Propi Super 25 EC/ Tilt 250E® (propiconazole)	3	2	Yes	30 days							X		
Quash® (metconazole)	3	2	Yes	21 days							X		X*
Serenade OPTI® (<i>Bacillus subtilis</i> QST 713 strain)	44	None	No	None						X			X
Topnotch® (azoxystrobin + propiconazole)	3 + 11	2	Yes	30 days	X		X				X		X*
Zolera™ FX (fluoxastrobin + tetraconazole)	3 + 11	2	No	14 days		X*	X*				X		X*

* Product registered for suppression only.

** Follow label for proper rotation of fungicides with different modes of action. Some products are not labelled for two sequential applications. Rates and staging may vary by fungicide and disease pressure. Read and follow label directions.

Version March 2021